



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$x^4 - (a^2 + b^2 + 2c^2)x^2 + 4abcx - c^2(a^2 + b^2 - c^2) = 0 \dots\dots\dots(5).$$

Restoring numbers in (5), we have

$$x^4 - 2518x^2 + 14400x - 22419 = 0.$$

Solving this equation by Horner's Method, we find $x = 47.145$ feet, nearly.

CALCULUS.

Conducted by J. M. COLAW, Monterey, Virginia. All contributions to this department should be sent to him.


SOLUTIONS OF PROBLEMS.

58. Proposed by O. W. ANTHONY, M. Sc., Professor of Mathematics, Columbian University, Washington, D. C.

A line passes through a fixed point and rotates uniformly about this point. Another line passes through a point which moves uniformly along the arc of a given curve and rotates uniformly about this point. Develop a method for finding the locus of intersection of these two lines. Apply to case of circle and straight line.

II. Solution by C. W. M. BLACK, A. M., Professor of Mathematics in Wesleyan Academy, Wilbraham, Massachusetts.

Let O be the origin, P_3 the fixed point, its coördinates being (r_3, θ_3) , and let AB be a given position of line through P_3 . Let $P_1(r_1, \theta_1)$ be position of point on curve and CD the line through it, both corresponding to the position AB of other line. Also let HK be position of AB revolved through an $\angle \psi$, and let $P_2(r_2, \theta_2)$ and EF be the corresponding position of P_1 and CD .



Let $r = f(\theta)$ be equation to curve P_1P_2 . Let η = the angle made by AB , and η_1 the one made by CD with a polar axis. Let a = angular rate of revolution of AB , and na of CD .

$\therefore \angle$ between CD and $EF = n\psi$.

Let b = linear rate of movement of P_1 . Then $\psi/a = P_1P_2/b \dots\dots\dots(1).$

Equation to KH is $r = [r_3 \sin(\eta + \psi - \theta_3)] / \sin(\eta + \psi - \theta) \dots\dots\dots(2).$

Equation to EF is $r = [r_2 \sin(\eta_1 + n\psi - \theta_2)] / \sin(\eta_1 + n\psi - \theta) \dots\dots\dots(3).$

By integration,

